

Sunyaev-Zeldovich Effect from Quasar-driven Blast-waves

P. Platania¹, C. Burigana², G. De Zotti³, E. Lazzaro⁴, and M. Bersanelli¹

¹*Università di Milano, Dipartimento di Fisica, Via Celoria 16, I-20133, Milano, ITALY*

²*Istituto TESRE, CNR, Via Gobetti 101, I-40129 Bologna, ITALY*

³*Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, ITALY*

⁴*Istituto di Fisica del Plasma, CNR, Via Cozzi 53, I-20125, Milano, ITALY*

Accepted ... 2002. Received ... 2002 ; in original form 2002 ...

ABSTRACT

Quasar-driven winds are currently the best candidates for accounting for the pre-heating of the intergalactic medium in clusters. Such winds, occurring during early phases of the evolution of spheroidal galaxies, shock-heat the interstellar gas, thus inducing a detectable Sunyaev-Zeldovich effect. We estimate the amplitude and the angular scale of such effect as well as its counts as a function of the Comptonization parameter y . The contamination due to radio emission by the quasar itself is also discussed. The corresponding mean Compton distortion of the cosmic microwave background spectrum is found to be well below the COBE/FIRAS upper limit.

Key words: quasars: general – galaxies: formation – cosmic microwave background

1 INTRODUCTION

It is natural to expect that extremely powerful sources, such as quasars, strongly affect the surrounding medium. The recent evidences of a tight relationship between black hole masses and velocity dispersions of the host galactic bulges (Ferrarese & Merritt 2000; Gebhardt et al. 2000a,b; Merritt & Ferrarese 2001a,b; McLure & Dunlop 2001) have strongly strengthened the case for a close connection between the evolutionary pathways of spheroidal galaxies and of quasars. Super-massive black holes have been found to be ubiquitous at the centers of local spheroidal galaxies (Kormendy & Richstone 1995; Magorrian et al. 1998; van der Marel 1999). The observed correlation between the mass of the black hole and that of the galaxy spheroidal component hints at a substantial feedback of the nuclear activity to the surrounding medium (Silk & Rees 1998; Monaco et al. 2000; Granato et al. 2001): energetic quasar-driven winds can sweep up the interstellar gas and halt both star formation and the growth of the central black hole.

If a significant fraction of the huge amount of energy released by quasars goes into ionization and heating of the neighboring gas, Compton cooling may produce a detectable Sunyaev-Zeldovich (1972; SZ) effect. The idea of strong shock heating of the medium originated by energetic outflows from early quasars, originally developed by Ikeuchi (1981), was recently revived by Natarajan, Sigurdsson & Silk (1998), Natarajan & Sigurdsson (1999), and Aghanim, Baland & Silk (2000). Evidences of strong quasar-driven winds can be seen in Broad Absorption Lines (BAL) quasars, comprising 10%–15% of optically selected quasars (Hamann & Ferland 1999 and reference therein). The dynamic interac-

tion of such energetic outflows with the surrounding protogalactic gas can heat it to high temperature. A release of an amount of mechanical energy from Active Galactic Nuclei several times larger than that produced by supernovae in the surrounding galaxies may be necessary to account for the pre-heating of the intergalactic medium in clusters (Wu et al. 2000; Kravtsov & Yepes 2000; Balogh et al. 2001; Bower et al. 2001). Natarajan & Sigurdsson (1999) suggested that the SZ effect associated to very energetic, quasar driven, winds may account for the reported isolated Cosmic Microwave Background (CMB) temperature decrements in directions where no clusters of galaxies are detected, but quasar pairs are present (Jones et al. 1997; Richards et al. 1997). Detection of this effect would obviously be informative on the physics of quasar/galaxy evolution.

In this paper we present a new investigation of the problem, taking explicitly into account the relevant energetics as well as the observed relationships between black hole and host galaxy properties. Our approach, focussing on energetics, is less liable to uncertainties ensuing from the poor knowledge of the details of the gas heating process. In Section 2 we introduce the basic ingredients entering our calculations and estimate the amplitude and the angular scale of the SZ effect. In Section 3 we present tentative estimates of its number counts. The results are discussed in Section 4.

2 SZ EFFECT FROM QUASARS

The main factors determining the amplitude of the quasar-driven SZ effect, usually measured in terms of the Comptonization parameter

$$y = \int_{-R}^R \frac{kT_e(r)}{m_e c^2} \sigma_T n_e(r) dr, \quad (1)$$

(where T_e , n_e and m_e are the electron temperature, density and mass, σ_T is the Thomson cross section, and R is the radius of the ionized region) can be illustrated as follows.

The fractional amount of energy, $\Delta\epsilon/\epsilon_{\text{CMB}}$, transferred to the CMB by Compton cooling of the plasma is related to y by (Zeldovich & Sunyaev 1969):

$$y \simeq \frac{1}{4} \frac{\Delta\epsilon}{\epsilon_{\text{CMB}}}, \quad (2)$$

where $\epsilon_{\text{CMB}} = aT_{\text{CMB}}^4 \simeq 4.2 \times 10^{-13} (1+z)^4 \text{ erg cm}^{-3}$.

In the redshift range where quasars are observed ($z \lesssim 6.3$, Fan et al. 2001b) and for realistic values of the density of the medium, energetic cosmological blast waves remain adiabatic to relatively low z , i.e. the cooling time is longer than the expansion timescale, t_{exp} (Voit 1996; see also the discussion below). Thus, if a fraction f_h of the total energy released by the quasar, E_{tot} , goes into heating of the gas, the amount of energy per unit volume transferred to the CMB through comptonization in a time $t < t_c$ is:

$$\Delta\epsilon \simeq \frac{f_h E_{\text{tot}}}{V} \frac{t}{t_c}, \quad (3)$$

where V is the volume occupied by the hot gas and $t_c = 3m_e c / 4\sigma_T aT_{\text{CMB}}^4 \simeq 7.3 \times 10^{19} (1+z)^{-4} \text{ s}$ is the Compton cooling timescale. We adopt, for simplicity, an Einstein-de Sitter cosmology, so that the expansion timescale is $t_{\text{exp}} = a(t)/\dot{a}(t) = (1/H_0)(1+z)^{-3/2}$, where $a(t)$ is the cosmic scale factor and $1/H_0 \simeq 6.17 \times 10^{17} h_{50}^{-1} \text{ s}$, $h_{50} = H_0/50 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2.1 Evolution of the shock within the quasar host galaxy

Close relationships between the mass of the central black hole, M_{bh} , and the properties of host elliptical galaxies or of galaxy bulges have been reported. Kormendy & Richstone (1995), Magorrian et al. (1998), and McLure & Dunlop (2001) found that M_{bh} is proportional to the luminosity of the spheroidal galaxy component. A tighter correlation between M_{bh} and the line-of-sight velocity dispersion, σ , was discovered by Gebhardt et al. (2000) and Ferrarese & Merritt (2000), although the two groups find somewhat different slopes. We adopt the relationship obtained by Gebhardt et al. (2000), based on a larger sample:

$$M_{\text{bh}} = 1.2 \cdot 10^8 \left(\frac{\sigma}{200 \text{ km/s}} \right)^{15/4} M_{\odot}. \quad (4)$$

The total amount of energy released by the quasar is $E_{\text{tot}} = \epsilon_R M_{\text{bh}} c^2$, where $\epsilon_R \simeq 0.1$ is the mass-to-energy conversion efficiency of the black hole.

We model the host galaxy as an isothermal sphere with density profile

$$\rho(r) = \rho_0 r^{-2}, \quad (5)$$

with

$$\rho_0 = \frac{\sigma^2}{2\pi G}, \quad (6)$$

and an outer cut-off radius R_g defined by the condition that the mean density within R_g is $\delta \simeq 200$ times higher than

the mean density of the universe at the redshift z_f when the galaxy formed [numerical simulations (Cole & Lacey 1996) show that this radius approximately separates the virialized and infall regions; see also Navarro et al. 1997]:

$$R_g = \frac{2\sigma}{H_0 \delta^{1/2} (1+z_f)^{3/2}} \simeq 130 h_{50}^{-1} \left(\frac{E_{\text{tot}}}{10^{62}} \frac{0.1}{\epsilon_R} \right)^{4/15} \left(\frac{\delta}{200} \right)^{-1/2} \left(\frac{3.5}{1+z} \right)^{3/2} \text{ kpc}. \quad (7)$$

Several lines of evidence indicate that massive ellipticals were already in place by $z \simeq 2.5$ (Renzini & Cimatti 1999; Ferguson et al. 2000; Daddi et al. 2000; Cohen 2001), although the issue is still somewhat controversial. For $z_f \simeq 2.5$ Eq. (7) yields for R_g values consistent with current estimates for massive galaxies (see, e.g., Peebles 1993). We assume that the distribution of baryons reflects that of dark matter.

It is interesting to note, in passing, that this model, together with the Gebhardt et al. (2000) relationship [Eq. (4)] implies $M_{\text{bh}} \propto M_{\text{DM}}^{1.25}$, M_{DM} being the total mass of the galaxy, dominated by the dark matter halo. McLure & Dunlop (2001) find a best-fit relationship between black-hole mass and R -band luminosity of the host galaxy bulge $M_{\text{bh}} \propto L_R^{1.525}$. Adopting, as they do, the relationship $M_{\text{bulge}} \propto L^{1.31}$, determined by Jørgensen et al. (1996), we have $M_{\text{bh}} \propto M_{\text{bulge}}^{1.16}$. If we adopt the Magorrian et al. (1998) dependence, $M_{\text{bulge}} \propto L^{1.18}$, which according to Laor (2001) may be more appropriate, we get $M_{\text{bh}} \propto M_{\text{bulge}}^{1.29}$. In both cases the observationally derived relationship between M_{bh} and M_{bulge} is very close to the one yielded by the present model if $M_{\text{bulge}} \propto M_{\text{DM}}$.

The radius R_s of a self-similar blast-wave carrying an energy $E_b = f_h E_{\text{tot}}$, expanding in a medium whose density varies as r^{-2} , increases with time as (Ostriker & McKee 1988):

$$R_s = \left[\frac{\xi E_b}{3\rho_0} \right]^{1/3} t^{2/3}, \quad (8)$$

where $\xi \approx 1.5$.

Using Eqs. (7) and (8) it is easily checked that, for realistic values of the parameters, the time, t_g , required for the shock front to reach the outer radius of the galaxy

$$t_g \simeq 8.9 \times 10^{16} h_{50}^{-3/2} \left(\frac{E_{\text{tot}}}{10^{62}} \right)^{1/6} \left(\frac{\epsilon_R}{0.1} \right)^{-2/3} \cdot \left(\frac{f_h}{0.1} \right)^{-1/2} \left(\frac{\delta}{200} \right)^{-3/4} (1+z)^{-9/4} \text{ s} \quad (9)$$

is always shorter than the expansion timescale t_{exp} . Therefore the shocks propagate outside of the host galaxy, and can heat up the general intra-cluster medium (ICM). However, due to the much lower electron density in the ICM, compared to the mean density within the galaxy, most of the SZ signal comes from within R_g (see also da Silva et al. 2001).

The main radiative cooling process for the redshifts of interest here ($z \leq 6$) is free-free, whose cooling timescale is:

$$t_{\text{ff}} = \frac{3n_e kT_e}{w_{\text{ff}}}, \quad (10)$$

where $w_{\text{ff}} = 1.4 \cdot 10^{-27} T_e^{1/2} n_e^2 \bar{g} \text{ erg s}^{-1}$ is the cooling rate of a plasma with electron number density n_e and temperature T_e . The present mean ratio between baryon (mostly in stars)

and dark matter mass in massive spheroidal galaxies is estimated to be $M_b/M_{DM} \simeq 0.03$ (McKay et al. 2001; Marinoni & Hudson 2002). An upper limit to the cooling rate (and, correspondingly, a lower limit to t_{ff}) is obtained assuming that essentially all such baryons were in the interstellar gas, so that

$$n_e(r) = 3 \times 10^{-2} \frac{\sigma^2}{2\pi G m_p r^2}, \quad (11)$$

where m_p is the proton mass, we find (setting $\bar{g} = 1$) that $t_{ff} < t_g$ [Eq. (9)] for

$$r < r_{ff} \simeq 35 h_{50}^{-3/4} \left(\frac{T_e}{10^6 \text{K}} \right)^{-1/4} \left(\frac{E_{\text{tot}}}{10^{62}} \right)^{7/20} \left(\frac{\epsilon_R}{0.1} \right)^{-3/5} \cdot \left(\frac{f_h}{0.1} \right)^{-1/4} \left(\frac{\delta}{200} \right)^{-3/8} (1+z)^{-9/8} \text{ kpc}, \quad (12)$$

i.e. for only a very small fraction of the volume heated by the blast wave [see Eq. (8)]. Therefore we neglect, in the following, the radiative losses.

2.2 Energetics of the shock

The fraction of power released by the quasars which goes into heating of the gas is highly uncertain. Natarajan & Sigurdsson (1999) assume this fraction to amount to about half of the bolometric luminosity. Analyses of the X-ray properties of the intra-cluster medium (ICM) and, in particular, of the temperature-luminosity relation (Ponman et al. 1999; Valageas & Silk 1999; Cavaliere et al. 2000; Tozzi & Norman 2001; Balogh et al. 2001) indicate the need of a substantial “pre-heating” of the gas before it collapses into the cluster. As discussed by Cavaliere & Menci (2001), measurements of the associated SZ effect very effectively probe the processes responsible for non-gravitational heating of the ICM. Bower et al. (2001) have shown that, unless the transfer of supernova energy to the ICM is very (perhaps unrealistically) efficient, additional heating sources, such as mechanical energy from quasar-driven winds are required. Adopting a baryon mass corresponding to $\Omega_b = 0.025 h^{-3/2}$ ($h = H_0/100$) and a fraction of baryons converted into stars of $f_{\text{gal}} = 0.16 h^{1/2}$, Bower et al. (2001) estimate that an energy of $\epsilon_{SN} \times 10^{49}$ erg per $M_\odot$ of stars formed, with $\epsilon_{SN} \simeq 1.3 - 2 h^{-1/2} = 1.8 - 2.8 h_{50}^{-1/2}$ must go into heating of the ICM, and that most of it is unlikely to come from supernova explosions. Other energy sources, the most obvious being quasar winds, must be advocated.

Adopting a ratio of $f_\star = 0.03$ between the mass in stars and the gravitational mass of spheroidal galaxies we have:

$$f_h \simeq 4.7 \times 10^{-2} \frac{f_\star}{0.03} \frac{\epsilon_{SN}}{2.8} \left(\frac{\epsilon_R}{0.1} \right)^{-4/5} \cdot \left(\frac{E_{\text{tot}}}{10^{62}} \right)^{-1/5} h_{50}^{-3/2} \left(\frac{\delta}{200} \right)^{-1/2} \left(\frac{1+z}{3.5} \right)^{-3/2}, \quad (13)$$

indicating that the mechanical power is a minor fraction of the bolometric luminosity.

2.3 Amplitude and angular scale of the SZ effect

As argued in sub-section 2.1, most of the SZ signal is expected to come from within the galaxy, so that we will set, in Eq. (3), $V = V_g = (4\pi/3)R_g^3$ and $t = t_g$. On the other




Figure 1. Differential counts per deg^2 and per unit interval of $\log(y)$ (the two upper curves) or of $\log(y_{\text{obs}})$ (the two lower curves) of SZ signals due to quasar-driven blast waves, as a function of the comptonization parameter y (or y_{obs}), for the reference values of the parameters. The two solid lines correspond to $t_q = 10^7$ yr, while the two dashed curves correspond to $t_q = 10^8$ yr. The two lower curves take into account the dilution effect of a Gaussian observing beam with $\text{FWHM} = 10'$.

hand, only a minor fraction f_g of the energy carried by the blast-wave goes into heating of the gas within the galaxy (most of it is dissipated in the ICM). Inserting in Eq. (2) the above results, using for f_h the expression of Eq. (13), and taking into account that, in the Rayleigh-Jeans region $(\Delta T/T)_{\text{RJ}} = -2y$ we end up with

$$\left| \left(\frac{\Delta T}{T} \right)_{\text{RJ}} \right| \simeq 2.2 \times 10^{-4} \frac{f_g}{0.1} \left(\frac{\epsilon_{SN}}{2.8} \right)^{1/2} \left(\frac{\epsilon_R}{0.1} \right)^{-4/15} \cdot \left(\frac{E_{\text{tot}}}{10^{62}} \right)^{4/15} \left(\frac{\delta}{200} \right)^{1/2} h_{50}^{3/4} \left(\frac{f_\star}{0.03} \right)^{1/2} \left(\frac{1+z}{3.5} \right)^{3/2}. \quad (14)$$

The angular radius is $\theta_{SZ} \simeq R_g/d_A$,

$$d_A(z) = \frac{2c}{H_0} \frac{1+z - (1+z)^{1/2}}{(1+z)^2} \quad (15)$$

being the angular diameter distance. From Eq. (7) we have:

$$\theta_{SZ} \simeq 17'' \left(\frac{E_{\text{tot}}}{10^{62}} \right)^{4/15} \left(\frac{\epsilon_R}{0.1} \right)^{-4/15} \left(\frac{1+z}{3.5} \right)^{-3/2} \frac{d_A(2.5)}{d_A(z)}, \quad (16)$$

where $d_A(2.5) = 1595/h_{50}$ Mpc. The amplitude and the angular diameter of the effect, $2\theta_{SZ} \simeq 34''$, for the reference values of the parameters, are not far from the results reported by Richards et al. (1997): $|\delta T/T| \sim 10^{-4}$ over an area of $30'' \times 65''$, and by Jones et al. (1997): $|\delta T/T| \sim 1.4 \times 10^{-4}$ in a beam of $100'' \times 175''$. Since powerful high- z quasars are highly clustered (Croom et al. 2001), the larger than expected observed angular scale might be interpreted as due to the combination of blast waves from neighbor quasars. It should be stressed, however, that in neither case the explanation in terms of quasar-driven blast waves is clearly supported by observations. A more conservative explanation is that the observed signals come from the SZ effect in previously unknown clusters.

3 COUNTS OF SZ SIGNALS FROM QUASARS

The epoch-dependent “luminosity function” of SZ signals, $\phi_{SZ}(y, z)$ can be roughly estimated from the luminosity function of quasars, given the quasar lifetime t_q . We have adopted one of the analytical evolutionary models for the B-band luminosity function, $\phi_B(L_B, z)d\log L_B dz$, proposed by Pei (1995): the two-power-law model with $h_{50} = 1$, $q_0 = 0.5$, and optical spectral index $\alpha_o = 0.5$ ($f(\nu) \propto \nu^{-\alpha}$). This model works well up to $z \simeq 4.5$, but appears to underpredict by a large factor the surface density of higher redshift quasars, which, however, are so rare (Fan et al. 2001a) that their contributions to the counts of the SZ signal is small.

The B-band luminosity is related to the total energy released by $E_{\text{tot}} = k_B L_B t_q$, where k_B is the bolometric correction for which we adopt the value $k_B = 6$ (the median value for the sample of Elvis et al. 1994, corrected to account for the different definition of L_B used by Pei 1995). If t_q is independent of both luminosity and redshift, and considering only the SZ effect within the host galaxy, we have:

$$\phi_{SZ}(y, z) = \frac{15}{4} \phi_B(L_B, z) \frac{t_q}{t_q}, \quad (17)$$

where $\phi_{SZ}(y, z)$ is the “luminosity function” of the SZ signal per unit $\log y$ and z intervals, and the factor $15/4$ comes from $d\log L_B/d\log y = d\log E_{\text{tot}}/d\log y$ [Eq. (14)]. The quasar lifetime is still very uncertain. Recent estimates (Salucci et al. 1999; Monaco et al. 2000; Martini & Weinberg 2001) suggest t_q to be in the range $8 \times 10^6 - 10^8$ yr, values $\simeq \text{few} \times 10^7$ yr being favoured (Haehnelt et al. 1998; Pentericci et al. 2002; Ciotti et al. 2001).

Inserting Eq. (13) in the expression for t_g [Eq.(9)] we have:

$$t_g \simeq 7.7 \times 10^{15} h_{50}^{-3/4} \left(\frac{E_{\text{tot}}}{10^{62}} \right)^{4/15} \left(\frac{\epsilon_R}{0.1} \right)^{-4/15} \cdot \left(\frac{\epsilon_{SN}}{2.8} \right)^{-1/2} \left(\frac{f_*}{0.03} \right)^{-1/2} \left(\frac{\delta}{200} \right)^{-1/2} \left(\frac{1+z}{3.5} \right)^{-3/2} \text{ s}. \quad (18)$$

Although the amplitude of the SZ effect is distance independent, the observed signal is affected by distance dependent beam dilution or resolution effects, so that the SZ signal observed with an instrumental solid angle ω_{beam} is:

$$y_{\text{obs}} \simeq y F(\omega_{\text{beam}}/\omega_{SZ}), \quad (19)$$

where $\omega_{SZ} = \pi \theta_{SZ}^2$ and

$$F(\omega_{\text{beam}}/\omega_{SZ}) \simeq \begin{cases} \omega_{\text{beam}}/\omega_{SZ} & \text{if } \omega_{\text{beam}} < \omega_{SZ} \\ \omega_{SZ}/\omega_{\text{beam}} & \text{if } \omega_{\text{beam}} > \omega_{SZ} \end{cases}. \quad (20)$$

For given z , $F \propto E_{\text{tot}}^{\pm 8/15} \propto y^{\pm 2}$, the sign of the exponent depending on the ratio $\omega_{\text{beam}}/\omega_{SZ}$ [see Eq. (20)], so that $y_{\text{obs}} \propto y^{\pm 2}$. The differential counts per steradian of y_{obs} then write:

$$\frac{dN_{SZ}}{dy_{\text{obs}}} = \frac{1}{\ln(10) y_{\text{obs}}} \int dz \frac{dV}{dz} \phi_{SZ}(y_{\text{obs}}/F, z) \frac{d\log y}{d\log y_{\text{obs}}}, \quad (21)$$

where, for a Friedman universe, the volume element per unit solid angle and unit dz is $dV/dz = (c/H_0) d_L^2 (1+z)^{-13/2}$, $d_L = (2c/H_0)(1+z - \sqrt{1+z})$ being the luminosity distance. The differential counts $dN_{SZ}/d\log y = y \ln(10) dN_{SZ}/dy$ are shown in Fig. 1 for two values of t_q , 10^7 yr and 10^8 yr. For both cases, the effect of dilution by a Gaussian beam with FWHM = $10'$ is also shown.

The extremely steep slope of the bright end of the counts is model dependent. It comes out from the combined effect of the steep slope of the high luminosity portion of the quasar luminosity function and of the mild dependence of y on E_{tot} [see Eq. (14)], implied by the present model. Clearly, the assumption of dispersion-less relationships between the quasar blue luminosity and E_{tot} , and between E_{tot} and y is an over-simplification. Dispersions will result in a flattening of the bright end of the counts.

The surface density of SZ signals is strongly dependent on the amount of thermal energy injected in the interstellar gas of the host galaxy, which, according to the present model, is proportional to $t_q^{4/5}$. For t_q in the range $10^7 - 10^8$ yr, we expect some 10^3 SZ signals per square degree with $y > 2 \times 10^{-4}$ to be detected with an imaging survey with $\lesssim 30''$ resolution (FWHM). For higher values of y , the expected surface density is strongly dependent on the quasar lifetime t_q . The survey should be carried out at $10 \text{ GHz} < \nu < 150 \text{ GHz}$ to avoid blurring of the SZ signal by local radio or dust emission. The signal is strongly diluted in most current surveys aimed at mapping CMB anisotropies with angular resolution $\gtrsim 5' - 10'$. Even high sensitivity, all sky surveys such as those to be carried out by ESA’s PLANCK mission, cannot efficiently detect such signals. Much better prospects for detecting SZ effects from quasar-driven blast waves are offered by ground based bolometric arrays and interferometers with arc-min or sub-arcmin resolution, such as the Arcminute MicroKelving Imager (Kneissl et al. 2001), and AMiBA (Lo et al. 2001).

Since, in the present framework, quasars are signposts of massive galaxies at high redshifts and are therefore strongly biased tracers of the matter distribution, we expect them (and the associated SZ signals) to be highly clustered.

4 DISCUSSION

4.1 Effect of local radio emission

It is entirely plausible, and perhaps required by evidences of a substantial preheating of the intra-cluster gas, that quasars inject in the surrounding medium an amount of energy sufficient to produce a detectable SZ effect. The corresponding angular scale is expected to be relatively small (sub-arcmin), so that the blurring effect by the emission from the quasar itself or from the host galaxy may be important.

To estimate to what extent the radio emission associated with the quasar might blur the SZ signal, we refer to the median ratio of monochromatic luminosities at 5 GHz and at $\nu_B \simeq 6.82 \times 10^{14}$ Hz (corresponding to $\lambda_B = 0.44 \mu\text{m}$), for the radio-loud and radio-quiet quasars in Table 2 of Elvis et al. (1994). We find $\log(l_{5\text{GHz}}/l_B)_{\text{median}} = -0.47$ for radio-quiet quasars and $= 2.2$ for radio loud.

Taking into account that the contribution to the antenna temperature at the frequency ν within a solid angle $\omega_{SZ} = \pi \theta_{SZ}^2$, of a source of 5 GHz flux $S_{5\text{GHz}}$ and spectral index $\alpha \simeq 0.7$ ($S_\nu \propto \nu^{-\alpha}$) is:

$$\Delta T_A = \frac{S_{5\text{GHz}} (\nu/5 \text{ GHz})^{-\alpha} c^2}{2k_b \nu^2 \omega_{SZ}}, \quad (22)$$

we have, for radio-quiet quasars, in the Rayleigh-Jeans re-

gion

$$\frac{\Delta T_A}{T_A} \simeq 1.1 \cdot 10^{-3} \left(\frac{E_{\text{tot}}}{10^{62}} \right)^{7/15} \left(\frac{\epsilon_R}{0.1} \right)^{8/15} h_{50}^2 \cdot \frac{10^7 \text{ yr}}{t_q} \left(\frac{5 \text{ GHz}}{\nu} \right)^{2+\alpha} \left(\frac{1+z}{3.5} \right)^{-\alpha}, \quad (23)$$

In the case of radio loud quasars the coefficient would be about 500 times higher and therefore the radio emission would easily overwhelm the SZ signal, except, perhaps, at high frequencies ($\simeq 100$ GHz). However, only a minor fraction ($\lesssim 10\%$) of quasars are radio loud. But even in the case of radio quiet quasars the radio emission may, at least partially, fill up the SZ dip at $\nu \lesssim 10$ GHz, particularly if the quasar is surrounded by an intense starburst, which is also radio bright, as is frequently the case at high z (Omout et al. 2001). The contamination by radio emission sinks down rapidly with increasing frequency, but at $\nu \gtrsim 150$ GHz, dust emission powered by star formation in the host galaxy, may take over, due to its spectrum steeply rising with increasing frequency.

On the other hand, the quasar lifetime and the duration of intense starbursts are generally shorter than the time t_q for the blast-wave to reach the boundary of the host galaxy. Therefore the SZ signals may be observed when the quasars that originated them are dead.

4.2 Global comptonization distortion

The mean distortion of the CMB spectrum due to comptonization by the electrons heated by the blast waves, measured by the mean value $\langle y \rangle$ of the parameter y integrated along the line of sight, is 1/4 of the fractional amount of energy injected in the CMB per unit volume:

$$\langle y \rangle = \frac{f_{\text{h,eff}}}{4} \int \int_0^\infty dz d \log L_B \frac{\phi_B(L_B, z) k_B L_B}{\epsilon_{\text{CMB}}(z)} \frac{t_{\text{exp}}}{t_c} \frac{dt}{dz}. \quad (24)$$

where $f_{\text{h,eff}}$ is the effective fraction of the total energy released by the quasars carried by blast waves. From the above discussion it follows that pre-heating of the ICM requires $f_{\text{h,eff}} \lesssim 0.1$. Using the double-power-law quasar luminosity function by Pei (1995), we find:

$$\langle y \rangle \simeq 2.4 \times 10^{-6} \frac{f_{\text{h,eff}}}{0.1}. \quad (25)$$

Using Pei's (1995) exponential luminosity function, the value of $\langle y \rangle$ increases by 20%. On the other hand, it may be noted that Eq. (24) includes the contribution (which turns out to be quite substantial) of low- z quasars to the integral (the contribution from the redshift range 0–2 is comparable to that from $2 \leq z \leq 4$); however, in the present framework, the most energetic blast-waves should be associated to early phases of the quasar/galaxy evolution. Thus the expected global comptonization distortion of the CMB spectrum induced by quasar driven blast waves is well below the COBE/FIRAS limit $\langle y \rangle < 1.5 \times 10^{-5}$ (Fixsen et al. 1996). This limit could be (marginally) exceeded only in the unrealistic case that the total energy emitted by all quasars over the entire life of the universe went into heating of the surrounding gas. This would also entail an excessive (by almost one order of magnitude) pre-heating of the ICM.

ACKNOWLEDGMENTS

Work supported in part by MIUR and ASI.

REFERENCES

- Aghanim N., Balland C., Silk, J., 2000, A&A, 357, 1
 Balogh M.L, Pearce F., Bower R.G., Kay S., 2001, MNRAS, 326, 1228
 Bower R.G., Benson A.J., Lacey C.G., Baugh C.M., Cole S., Frenk C.S., 2001, MNRAS, 325, 497
 Cavaliere A., Giacconi R., Menci N., 2000, ApJ, 528, 77
 Cavaliere A., Menci N., 2001, MNRAS, 327, 488
 Ciotti L., Haiman Z., Ostriker J.P., 2001, proc. ESO symp. "The Mass of Galaxies at Low and High Redshift", R. Bender & A. Renzini eds., in press, astro-ph/0112131
 Cohen J.G., 2001, AJ, 121, 2895
 Cole S., Lacey C., 1996, MNRAS, 281, 716
 Croom S.M., Shanks T., Boyle B.J., Smith R.J., Miller L., Loaring N.S., Hoyle F., 2001, MNRAS, 325, 483
 Daddi E., Cimatti A., Renzini A., 2000, A&A, 362, L45
 da Silva A.C., Kay S.T., Liddle A.R., Thomas P.A., Pearce F.R., Barbosa D., 2001, ApJ, 561, L15
 Elvis M., Wilkes B.J., McDowell J.C., Green R.F., Bechtold J., Willner S.P., Oey M.S., Polomski E., Cutri R., 1994, ApJS, 95, 1
 Fan X., Strauss M.A., Schneider D.P., et al., 2001a, AJ, 121, 54
 Fan X., Narayanan V.K., Lupton R.H., et al., 2001b, AJ, 122, 2833 astro-ph/0108063
 Ferguson H.C., Dickinson M., Williams R., 2000, ARA&A, 38, 667
 Ferrarese L., Merritt, 2000, ApJ, 539, L9
 Fixsen D.J., Cheng E.S., Gales J.M., Mather J.C., Shafer R.A., Wright E.L., 1996, ApJ, 473, 576
 Gebhardt K., Bender R., Bower G., et al., 2000a, ApJ, 539, L13
 Gebhardt K., Kormendy J., Ho L.C., et al., 2000b, ApJ, 543, L5
 Granato G.L., Silva L., Monaco P., Panuzzo P., Salucci P., De Zotti G., Danese L., 2001, MNRAS, 324, 757
 Hamann F., Ferland G., 1999, ARA&A, 37, 487
 Haehnelt M.G., Natarajan P., Rees M.J., 1998, MNRAS, 300, 827
 Ikeuchi S., 1981, PASJ, 33, 211
 Jones M.E., Saunders R., Baker J.C., Cotter G., Edge A., Grainge K., Haynes T., Lasenby A., Pooley G., Röttgering H., 1997, ApJ, 479, L1
 Jørgensen I., Franx M., Kjaergaard P., 1996, 280, 167
 Kneissl R., Jones M.E., Saunders R., Eke V.R., Lasenby A.N., Grainge K., Cotter G., 2001, MNRAS, 328, 783
 Kormendy J., Richstone D., 1995, ARA&A, 33, 581
 Kravtsov A.V., Yepes G., 2000, MNRAS, 318, 227
 Laor A., 2001, ApJ, 553, 677
 Lo K.Y., Chiueh T.H., Martin R.N., et al., 2001, 20th Texas Symposium on Relativistic Astrophysics, 172
 Magorrian J., Tremaine S., Richstone D., Bender R., Bower G., Dressler A., Faber S.M., Gebhardt K., Green R., Grillmair C., Kormendy J., Lauer T., 1998, AJ, 115, 2285
 Marinoni C., Hudson M.J., 2002, ApJ, 569, 101
 Martini P., Weinberg D.H., 2001, ApJ, 547, 12
 McKay T.A., Sheldon E.S., Racusin J., et al., 2001, ApJ, submitted, astro-ph/0108013
 McLure R.J., Dunlop J.S., 2001, MNRAS, 327, 199
 Merritt D., Ferrarese L., 2001a, ApJ, 547, 140
 Merritt D., Ferrarese L., 2001b, MNRAS, 320, L30
 Monaco P., Salucci P., Danese L., 2000, MNRAS, 311, 279
 Natarajan P., Sigurdsson S., 1999, MNRAS, 302, 288
 Natarajan P., Sigurdsson S., Silk, J., 1998, MNRAS, 298, 577
 Navarro J.F., Frenk C.S., White S.D.M., 1997, ApJ, 490, 393
 Omout A., Cox P., Bertoldi F., McMahon R.G., Carilli C., Isaak K.G., 2001, A&A, 374, 371

- Ostriker J.P., McKee C.F., 1988, *Rev. of Mod. Phys.*, 60, 1
- Peebles P.J.E., 1993, *Principles of Physical Cosmology*, Princeton University Press, Princeton
- Pei Y., 1995, *ApJ*, 438, 623
- Pentericci L., Fan X., Rix H.W., Strauss M.A., Narayan V.K., Richards G.T., Schneider D.P., Krolik J., Heckman T., Brinkmann J., Lamb D.Q., Szokoly G.P., 2002, *AJ*, 123, 2151
- Ponman T.J., Cannon D.B., Navarro J.F., 1999, *Nature*, 397, 135
- Renzini A., Cimatti A., 1999, in *The Hy-Redshift Universe: Galaxy Formation and Evolution at High Redshift*, ASP Conf. Ser. 193, 312
- Richards E.A., Fomalont E.B., Kellermann K.I., Partridge R.B., Windhorst R.A., 1997, *AJ*, 113, 1475
- Salucci P., Szuskiewicz E., Monaco P., Danese L., 1999, *MNRAS*, 307, 637
- Silk J., Rees M.J., 1998, *A&A*, 331, L1
- Sunyaev R.A., Zeldovich Ya.B., 1972, *Comm. Ap. Sp. Phys.*, 4, 173
- Tozzi P., Norman C., 2001, *ApJ*, 546, 63
- Valageas P., Silk J., 1999, *A&A*, 350, 725
- van der Marel, R.P., 1999, *AJ*, 117, 744
- Voit M. G., 1996, *ApJ*, 145, 560
- Wu K.K.S., Fabian A.C., Nulsen P.E.J., 2000, *MNRAS*, 318, 889
- Zeldovich Ya.B., Sunyaev R.A., 1969, *Ap. Space Sci.*, 4, 301

This paper has been produced using the Royal Astronomical Society/Blackwell Science $\LaTeX$ style file.